

The Effect of Chinese Traditional Exercise-Baduanjin on Physical and Psychological Well-Being of College Students: A Randomized Controlled Trial.

Li M1, Fang Q1, Li J1, Zheng X1, Tao J1, Yan X2, Lin Q2, Lan X1, Chen B1, Zheng G1, Chen L3.

Author information

Abstract

BACKGROUND:

The physical and mental health of college students tends to continuously decline around the world, therefore, it is important to improve their health during college period. Baduanjin, a traditional Chinese exercise which combines movements with breath and mind, may be one of the selectable effective exercises. However, the effect of Baduanjin exercise on college students has not been established. In this study, we systematically assessed the effectiveness and safety of Baduanjin exercise on physical and mental health of college students by a rigorous randomized, parallel-controlled design.

METHODS:

A total of 222 college students from Fujian University of Traditional Chinese Medicine were recruited and randomly allocated at an equal ratio into control or Baduanjin training. Participants in control group were informed to maintain their original activity habit, and those in Baduanjin exercise group received a 12-week Baduanjin exercise training with a frequency of 1 hour per day and 5 days per week on the basis of their original activity habit. The physical and psychological outcomes, including lumbar muscle strength, lower limb proprioception function, physical fitness, as well as self-reported symptom intensity, stress, self-esteem, mood, quality of life, quality of sleep, and adverse events, were evaluated at baseline, 13 weeks (at the end of 12-week intervention), and 25 weeks (after the 12-week follow-up period). Intention-to-treat analysis was performed for the above outcomes.

RESULTS:

Compared with controls, significant improvements in Baduanjin exercise group at the end of 12-week intervention period were found on lower limb proprioception function (the rate of average trace error on right lower limb (%): control 23.50 ± 5.50 , Baduanjin 21.92 ± 6.54 , $P=0.004$; the rate of average trace error on left lower limb (%): control 22.32 ± 6.62 , Baduanjin 20.63 ± 4.62 , $P=0.046$), cardiorespiratory endurance (step test index: control 47.66 ± 5.94 , Baduanjin 50.07 ± 9.30 , $P=0.025$), flexibility (control

14.35±7.26cm, Baduanjin 15.39±6.43cm, P=0.009) and explosive force of lower limb (standing long jump test (m): control 1.77±0.24, Baduanjin 1.79±0.22, P=0.005 for adjustment baseline) in physical outcomes, and attention (Schulte Grid test (second): control 210.4±51.15, Baduanjin 192.4±47.14, P=0.034) in mental outcome. Lumbar muscle strength in Baduanjin group had been moderately enhanced but no significant difference compared to controls. No significant changes in other physical and mental outcomes, including vital capacity, blood pressure, heart rate, hand grip force, self-symptom intensity, stress, self-efficacy, quality of life, and quality of sleep, were found between groups. No adverse event was reported during the study period.

CONCLUSION:

Regular Baduanjin exercise had an advantage for college students on improvement of lower limb proprioception, enhance of cardiorespiratory endurance, flexibility, explosive force of lower limb and attention, compared with usual exercise.

The Effect of Chinese Traditional Exercise-Baduanjin on Physical and Psychological Well-Being of College Students: A Randomized Controlled Trial

[Moyi Li](#),¹ [Qianying Fang](#),¹ [Junzhe Li](#),¹ [Xin Zheng](#),¹ [Jing Tao](#),¹ [Xinghui Yan](#),² [Qiu Lin](#),² [Xiulu Lan](#),¹ [Bai Chen](#),¹ [Guohua Zheng](#),^{1,*} and [Lidian Chen](#)^{3,*}

Maciej Buchowski, Editor

[Author information](#) ► [Article notes](#) ► [Copyright and License information](#) ►

This article has been [cited by](#) other articles in PMC.

Go to:

Abstract

article-meta

Background

The physical and mental health of college students tends to continuously decline around the world, therefore, it is important to improve their health during college period. Baduanjin, a traditional Chinese exercise which combines movements with breath and mind, may be one of the selectable effective exercises. However, the effect of Baduanjin exercise

on college students has not been established. In this study, we systematically assessed the effectiveness and safety of Baduanjin exercise on physical and mental health of college students by a rigorous randomized, parallel-controlled design.

Methods

A total of 222 college students from Fujian University of Traditional Chinese Medicine were recruited and randomly allocated at an equal ratio into control or Baduanjin training. Participants in control group were informed to maintain their original activity habit, and those in Baduanjin exercise group received a 12-week Baduanjin exercise training with a frequency of 1 hour per day and 5 days per week on the basis of their original activity habit. The physical and psychological outcomes, including lumbar muscle strength, lower limb proprioception function, physical fitness, as well as self-reported symptom intensity, stress, self-esteem, mood, quality of life, quality of sleep, and adverse events, were evaluated at baseline, 13 weeks (at the end of 12-week intervention), and 25 weeks (after the 12-week follow-up period). Intention-to-treat analysis was performed for the above outcomes.

Results

Compared with controls, significant improvements in Baduanjin exercise group at the end of 12-week intervention period were found on lower limb proprioception function (the rate of average trace error on right lower limb (%): control 23.50 ± 5.50 , Baduanjin 21.92 ± 6.54 , $P=0.004$; the rate of average trace error on left lower limb (%): control 22.32 ± 6.62 , Baduanjin 20.63 ± 4.62 , $P=0.046$), cardiorespiratory endurance (step test index: control 47.66 ± 5.94 , Baduanjin 50.07 ± 9.30 , $P=0.025$), flexibility (control 14.35 ± 7.26 cm, Baduanjin 15.39 ± 6.43 cm, $P=0.009$) and explosive force of lower limb (standing long jump test (m): control 1.77 ± 0.24 , Baduanjin 1.79 ± 0.22 , $P=0.005$ for adjustment baseline) in physical outcomes, and attention (Schulte Grid test (second): control 210.4 ± 51.15 , Baduanjin 192.4 ± 47.14 , $P=0.034$) in mental outcome. Lumbar muscle strength in Baduanjin group had been moderately enhanced but no significant difference compared to controls. No significant changes in other physical and mental outcomes, including vital capacity, blood pressure, heart rate, hand grip force, self-symptom intensity, stress, self-efficacy, quality of life, and quality of sleep, were

found between groups. No adverse event was reported during the study period.

Conclusion

Regular Baduanjin exercise had an advantage for college students on improvement of lower limb proprioception, enhance of cardiorespiratory endurance, flexibility, explosive force of lower limb and attention, compared with usual exercise.

Trial Registration

Chinese Clinical Trial Registry ChiCTR-TRC-13003329 <http://www.chictr.org>

Go to:

Introduction

With the progress of science and technology, the integration of digital technology within daily life is becoming the cultural norm. More young people spend a majority of their spare time in the digital devices [1, 2]. College students are the vanguard of the first cohort of young people, and sedentary lifestyle becomes prevalent among college students globally [3–5]. A report from American College Health Association indicated that only 43.6% college students reported vigorous physical activity for 20 minutes or moderate activity for 30 minutes on only 3 or fewer days of the previous 7 days [6, 7]. Participation in large amounts of sedentary lifestyle is associated with multiple health problems such as impaired lipid profiles and glucose uptake [8]. As a result, college students may be at increased risk for the metabolic syndrome and susceptible to serious chronic diseases later in life [9–11]. Similarly, mental health problems in college students are increasing steeply due to peer pressure and environmental changes, and becoming a worldwide public health burden [12–14]. In the United States, mental disorders accounted for nearly half of the disease burden for young adults [15]. These mental health problems are associated with poorer academic achievements, instable intimate relationship, even suicide [16]. However, only a small part of college students with mental distress ask for and receive adequate attention [17]. It is well established that physical inactivity in young adults are directly linked to mental distress [18,19]. Therefore, physical activity may play an important role in the management of mild-to-moderate mental distress [20,21].

Regular physical activities or exercises have been demonstrated to benefit participants' physical, mental, and social well-being [22–24]. Recent studies showed that exercise for 8 to 12 weeks was effective for promoting physical fitness and mental health, and improving body composition in college students [25,26]. Baduanjin exercise, translated as the 'eight section of brocades', is one of many traditional Chinese Qigong exercises for health promotion. It includes eight separate, delicate, and smooth exercise movements, and each movement benefits different physical parts of body or particular organs [27]. Baduanjin exercise can promote Qi function through the whole exercise of body posture, movement, breathing, and meditation—that means using the natural energies to optimize and balance internal energy with the coordination of body, breath, and mind [28]. Current studies have suggested that Baduanjin training appeared to have substantive benefits for older adults [29–31]. A few studies for young adults also indicated that Baduanjin exercise provides potential benefits on reducing depression, stress and anxiety, building self-control and healthy mind, and enhancing physical function [32–34]. However, due to the methodological limitations in previous studies, whether Baduanjin exercise can be recommended as an effective exercise to enhance college students' health is still unknown. The purpose of this trial is to systematically evaluate the effect of Baduanjin exercise on physical and psychological outcomes of the college students in China.

Go to:

Methods and Design

2.1 Study design

This was a randomized, single-blind, parallel-controlled trial to evaluate the effectiveness and safety of Baduanjin exercise for the physical and mental health of college students. A total of 222 eligible college students from the Fujian University of Traditional Chinese Medicine (FJTCM) were recruited and randomly allocated to either control or Baduanjin exercise group with a ratio of 1:1. The relative physical and psychological outcomes were measured at baseline, 13 weeks (at the end of intervention), and 25 weeks (after the 12-week follow-up period). The design of the study was detailed in the published protocol [35].

2.2 Ethics approval

This trial was carried out in accordance with the declaration of Helsinki and approved by the Ethics Board of FJTCM (approval number: 2013, Reviewed-No. 043). All participants provided written informed consent prior to participation. The individual in this manuscript has given written informed consent (as outlined in PLOS consent form) to publish these case details.

2.3 Participants

Participants were college students recruited from the first or second grade in FJTCM through the campus poster, advertisement, and campus broadcast between September 9 and September 19, 2013.

Participants were eligible to participate in the study if they met the following criteria: aged 18 to 25 years; provided the informed consent; and were freshmen or sophomores. Participants were excluded if they had been engaged in a long-term regular practice of Baduanjin, or were members of the Martial Arts Association, Dance Association, Aerobics Association, Sanda Association, or Taekwondo Association of FJTCM; or if they had suffered from severe cardiovascular diseases, musculoskeletal system diseases, or sports contraindications. Baseline assessments were conducted within 2 weeks before the beginning of the intervention program.

2.4 Randomization, allocation concealment and blinding

The random allocation sequence was produced by an independent statistician who worked in the Evidence-Based Center of FJTCM via the PLAN sentences of the statistical software SAS9.1. The detail of random allocation sequence was kept with a project manager. After the baseline assessments, the project manager informed each eligible participant his/her group (Baduanjin exercise or control group) via cell phone.

Although it was impossible to blind the participants and exercise instructors, we masked participants' allocation information to outcome assessors, and urged participants not to disclose their allocation information during the period of outcomes measurement. In addition, we built the blind code of allocation to blind the statistician, and in which, the group assignment (Baduanjin exercise or usual exercise control group) was replaced by the alphabet A or B.

2.5 Intervention and follow-up

2.5.1 Control group

Participants in the control group were informed to keep their original physical activity habit during the 12-week intervention period.

2.5.2 Baduanjin exercise group

On the basis of their original physical activity habit, participants in the Baduanjin exercise group gathered and practiced 1 hour Baduanjin exercise at 5 p.m. each day with a frequency of 5 day per week at the gymnasiums of the university. Baduanjin exercise was instructed by two qualified instructors who had engaged in the physical education over 5 years. The training scheme of Baduanjin exercise was in accordance with the *Health Qigong—Baduanjin* published by the General Administration of Sport of China, which consisted of 10 postures (including the beginning and ending posture) [36].

Intervention period lasted 12 weeks from September 23, 2013 to December 15, 2013. All participants were required to fill in the daily activity log during the intervention period, in which the type, frequency and duration of participant's activity in a whole day was recorded and classified into low, moderate, or high intensity grade. An unsupervised 12-week follow-up period for each participant began on December 16, 2013, and ended on March 7, 2014. No participants received additional exercise intervention but were required to fill in the daily activity log during the 12-week follow-up period.

2.6 Outcome assessment

The primary measure of outcomes was the change in the lumbar muscle strength, lower limb proprioception function, as well as self-reported symptom intensity, stress, self-efficacy and attention measured at 13 weeks (the first week after completion of the 12-week intervention period). We also evaluated the long term effect over a 12-week period at 25 weeks (the first week after completion of 12-week follow-up period). Lumbar muscle strength (consisting of flexion, and rotation myodynamia) and lower limb proprioception function were assessed using the *Tergumed Work Station* (produced by *Proxomed GmbH, Germany*) and the *Prokin* proprioception evaluation and training system (produced by *Tecnobody.S.r.l, Italy.*) by the blinded operators at the Evaluation Department of Rehabilitation Hospital Affiliated to FJTCM.

The scores of self-reported symptom intensity, stress, self-efficacy and attention were felt to best reflect the wellbeing health condition of college students, and were conducted using their corresponding scales with the confirmed validity [37–39].

Secondary outcomes included changes in physical fitness, mood and mindfulness, self-esteem, quality of life, and quality of sleep measured at 13 weeks and 25 weeks. Physical fitness, consisting of cardiopulmonary function, hand grip force, explosive force of lower limb and flexibility, was measured using the step test, hand grip strength test, standing long jump and the ‘sit and reach’ test respectively, which has been shown to be a reliable and responsive measure in overall physical fitness of college students according to the *Chinese University Students’ Physical Health Standards*. The psychological health and life or sleep quality outcomes were collected via the self-reported questionnaires at the Center of College Student Activity of FJTCM. Details about the outcome assessment were described in the published protocol of this trial [35].

2.7 Sample size and statistical analysis

No previous study of Baduanjin exercise for the physical and mental well-being in college students with rigorous design was available. Although Baduanjin exercise as a whole may be broken down into eight separate movement form which each focuses on different physical area, majority of them need rotate the body on waist when practicing [40]. Therefore the change of lumbar muscle strength was established to estimate the sample size. Our preliminary data from 20 college students indicated the means with standard deviation of the flexion strength of lumbar muscle was 258.25 newton (N) and 114.19 N, respectively. The current trial was designed to detect a 20% increase on lumbar flexion muscle strength after the Baduanjin exercise intervention for 12-week period [41,42]. A sample size of 101 per group with a type I error of 5% ($\alpha = 0.05$) and 90% power ($\beta = 0.10$) was necessary. With an estimated dropout rate of 10%, a final sample size of 111 per group was required. The primary and secondary outcomes were analyzed on an intention-to-treat (ITT) basis. Method of Last Observation Carry Forward was applied to fill in the missing data. Analyses were done using SPSS 21.0 (IBM, Chicago, IL, USA) software packages. The statistical significance

was defined as two-sided P value of <0.05 .

In descriptive analysis of the sample, continuous variables were expressed as mean and standard deviation for normal distribution, and median and inter-quartile range for non-normal distribution. Normality was tested using Kolmogorov-Smirnov test. Appropriate transformations were applied in cases of non-normal distribution. Categorical variables were expressed as proportions with their standard error.

Baseline characteristics between groups were compared using the t -test or Mann-Whitney test for continuous variables and Pearson chi-squared or Fisher's exact test for categorical variables. If incomparability appears, the inequality factors were treated as confounding variables in the final efficacy analysis.

For comparison of the primary or secondary outcomes between groups, a t -test or non-parametric tests was used for continuous variables, and Pearson chi-squared or Fisher exact test for categorical variables. To control for possible confounding factors, linear regression model was applied for dependent continuous variables and logistic regression models for dependent categorical variables. Subgroup analysis stratified by gender was conducted for the primary outcomes.

Go to:

Results

3.1 Baseline characteristics

[Fig 1](#) illustrates the participants flow. 16 participants (10 in Baduanjin exercise group and 6 in control) dropped out before baseline assessment. Therefore they were excluded from the ITT analysis. During the 24-week study period (including 12-week intervention period and 12-week follow-up period), 5 participants dropped out and 7 participants lost to follow-up, with 93 (92.1%) participants in Baduanjin exercise group and 101 (96.2%) participants in control group. There was no statistically significant difference between groups in terms of drop-out or lost to follow-up ($\chi^2 = 1.586$, $P = 0.208$). Of 96 participants in the Baduanjin exercise group with the completed 12-week Baduanjin exercise training, 18 participants had 75% attendance rate (actual training days/plan training days) or less, and 38 participants possessed 76%-85%, 40 participants attained more than 85% attendance rate. Reasons given for not attending training mainly included training time clashing with other

commitments and illness.
 fig ft0fig mode=article fl



Fig 1

caption a4

caption a8 **Participant flow through the study reported in a CONSORT diagram.**

In 206 eligible participants, 82.5% of them were female students; the average age of participants was 20.78 years old. Baseline demographic characteristics between groups including sex proportion, the average of age, baseline exercise time, height and weight had no significant difference ([Table 1](#)). The average time of high intensity activities, low intensity activities, and sedentary time between groups were not statistically different whether in the intervention period or the follow-up period. The average time of moderate intensity activities in Baduanjin exercise group were higher than the control group during 12-week intervention period ($P = 0.001$) ([Table 2](#)).

table ft1table-wrap mode=article t1

Characteristics	Baduanjin exercise group (n = 103)	Control group (n = 103)
Total included in the final analysis	103	103
Sex: Female (%)	85 (82.1)	84 (80.6)
Age-yr (Mean(SD))	20.83±1.03	20.92±1.05
Height-cm (Mean(SD))	168.06±6.87	168.21±6.70
Weight-kg (Mean(SD))	53.62±7.41	53.96±8.03
Baseline exercise time* min	34.08±20.92	37.53±27.81

*Baseline exercise time was the self-reported mean time of undergone moderate-intensity exercise one month.

[Table 1](#)
caption
a4
caption
a8 **Baseli
ne
characte
ristics
between
Baduanji
n
exercise
and
control
groups.**

table ft1 table-wrap mode=article t1

Characteristics	Baduanjin exercise group (n = 48)	
Average activities time during the intervention period (1-12 weeks) :		
Sedentary time	7.77±2.83	∗
low intensity activities time	1.81±0.70	∗
moderate intensity activities time	1.06±0.35	∗
high intensity activities time	0.14±0.21	∗
Average activities time during the follow-up period (13-24 weeks) :		
Sedentary time	6.88±3.15	∗

[Tab
le 2](#)
cap
tion
a4

caption a8 **Comparison of average activities time during the 12-week intervention and 12-week follow-up period between groups (hours)**
3.2 The effect of Banduanjin exercise on primary physical and psychological outcomes

Although participants in Buduanjin exercise group had a mean improvement at the end of 12-week exercise of 42.24N, 52.13N, and 38.67N, respectively on flexion, left rotation, and right rotation, the difference was not statistically significant when compared with control group. The lower limb proprioception function in participants of Baduanjin exercise group had an obvious improvement after 12-week

regular exercise. The rate of average trace error (ATE) on right, and left lower limb decreased a mean of 7.29, 10.16 percent, respectively, while controls had a mean decline of 6.76, 9.59 percent respectively, for a significantly statistical difference ($P = 0.004$, $P = 0.046$, respectively). The primary psychological outcomes on well-being included self-reported psychological symptom intensity, stress, self-efficacy, and attention, which were assessed by using their corresponding questionnaires with the definite validity and reliability. No significant change in self-reported psychological symptom intensity, stress, self-efficacy was observed at the end of 12-week intervention period, respectively. However, the improvement of attention was significant, and participants with Baduanjin exercise improved a mean of 21.51 second over 12 weeks, while control group participants had a mean improvement of 14.24 second over 12 weeks after adjusting for the baseline values ($P = 0.034$) (Table 3). The above primary outcomes after additional 12-week follow-up were not different between groups (Table 3).

table ft1table-wrap mode=article t1

Outcomes	Group	Number of participants	Baseline	End of 12 weeks intervention period	Difference of 12 week intervention value to baseline	p^1
			Mean (SD)	Mean (SD)	Mean (SD)	
Lumbar muscle strength	BDJ	101	254.50 (109.32)	296.82 (112.64)	42.33 (71.81)	

		Table 3 caption a4
--	--	--------------------

caption a8 Primary physical and psychological effects. 3.3 The effect of Baduanjin exercise on secondary physical and psychological outcomes

On the physical parameter for the cardiorespiratory endurance was measured by using a method of step test, in which higher test index indicated better cardiorespiratory endurance. Compared to their baseline data, participants in control group had a decline of 5.62 point as well as a decline of 1.4 point in the Baduanjin exercise group at the end of 12-week intervention period with a significant difference between groups (P

= 0.025). Participants in Baduanjin exercise group had obvious improvement on physical parameters including flexibility, explosive force of lower limb compared with the controls at the end of 12-week intervention period ($P = 0.009$, $P = 0.005$, respectively) after adjusting for baseline values (Table 4). Other physical parameters, such as vital capacity, blood pressure, heart rate, and hand grip force, had no statistical difference at the end of intervention period between groups (Table 4). No significant change was observed at the end of 12-week intervention period or after 12-week follow-up period between groups for the other secondary psychological outcomes included self-esteem, mood and mindfulness, quality of life, and quality of sleep (Table 4).


Outcomes	group	numbers	Baseline	P value	End of 12 weeks	Difference of	p.
					value	12 week	
			Mean (SD)		Mean (SD)	to baseline	
Physical fitness							
Cardiorespiratory endurance (Step test index)	BDJ	101	56.47 (13.87)	0.328 (0.001)	50.87 (9.30)	-1.40 (16.42)	0.825
	Con	105	53.26 (11.08)	0.328 (0.001)	47.66 (8.94)	-5.62 (13.69)	0.825

				Table 4 caption a4
--	--	--	--	--------------------

caption a8Secondary physical and psychological effects.3.4
Subgroup analysis

Pre-specified subgroup analysis was performed to determine whether we could identify a subgroup most likely to benefit from the Baduanjin exercise on the primary outcomes. The left and right lumbar rotation strength in female participants with Baduanjin exercise had a moderate improvement comparing to female controls at end of additional 12-week follow-up period with a mean improvement of 56.92N and 45.36N versus 44.88N and 34.83N, respectively ($P = 0.066$, and $P = 0.042$) (Fig 2). Only attention in primary well-being health outcomes in female participants of Baduanjin exercise group had a significant improvement with a mean decline 21.38 second over intervention period comparing with a mean decline 10.83 second in controls ($P = 0.02$) (Fig 2).

fig ft0fig mode=article fl



				Fig_2 caption a4
--	--	--	--	--

caption a8Subgroup analysis.3.5 Safety

No adverse events (AEs) related to Baduanjin exercise occurred in this trial.

Go to:

Discussion

In this randomized controlled trial of Baduanjin exercise comparing with control for the physical and mental health of college students, we found significant improvement with regular Baduanjin exercise on lower limb proprioception function, cardiorespiratory endurance, flexibility, explosive force of lower limb, and attention over a 12-week intervention than usual exercises. However, significant benefits were not observed over the additional 12-week follow-up period, which indicated there was no long-term benefit to practice Baduanjin exercise. No adverse event related to Baduanjin exercise was reported during the intervention period, which suggested this form of exercise is safe.

As one of the most common forms of traditional Chinese qigong exercises, Baduanjin exercise is not only easy to learn, but also has less cognitive demanding [43]. It has been considered as a popular community exercise to promote health in China [44]. According to traditional Chinese medicine theory, good health means the unison and balance of the body and spirit. The benefits of regular Baduanjin exercise are expressed through adjusting breathing to make the process of smoother, unifying mind and breathing, strengthening muscles and tendons to make the body more flexible and the unison of mind and body [45,46]. In view of sport medicine and traditional Chinese medicine, practitioners should focus on their lumbar function, and it stresses ‘take the waist as the axis’ when practicing [47]. It is therefore expected to enhance participants’ lumbar muscle strength. In this trial, the increments of lumbar muscle strength in college students for 12

weeks of regular Baduanjin exercise are higher than those with usual exercise, however, the statistical significant difference was not observed between groups. The explanation may be that the college students in control group were already involved in high-intensity sport activities such as basketball, badminton, and ping-pong, etc. However, the enhance on lower limb proprioception function after 12 weeks of Baduanjin exercise in college students was higher than controls. Studies in community-dwelling older population indicated that a Baduanjin exercise program could modulate blood lipid metabolism, insulin sensitivity, and sleep quality [29–31,48]. However, there are few studies focusing on its effect among young adults, especially college students. A previous study with 24 enrolled college students demonstrated an improvement on blood pressure, heart rate, lung capacity, weight, and hand grip strength after 12-week Baduanjin exercise [49]. The significant changes on vital capacity and heart rate were reported in another study with 200 college students [50]. Both studies were not randomized controlled trials, therefore it is difficult to generalize the study findings. In this randomized parallel controlled study comparing with usual exercise, the significant changes of vital capacity, blood pressure, heart rate, and hand grip force between groups were not observed. However, the effect of Baduanjin exercise on cardiorespiratory endurance, flexibility and explosive force of lower limb had obvious improvement over 12-week regular practice. Baduanjin exercise can be described as a combination of Confucianism, Buddhism, Taoism and traditional Chinese medicine. The key point when practicing is thought to be regulation of the mind. Therefore, it requires practitioner to remove all thoughts and focus on a certain region of the body to feel the flow of Qi through the body [51]. Through the practice of Baduanjin exercise, one can expect to improve, strengthen, and gain spiritual cultivation. A previous study found that regular Baduanjin exercise with 3 times per week and 60 min per time for 8 weeks had a positive trend to improve self-image and reduce psychologic distress comparing with the inactive controls [52]. Another study from female university students suggested that Baduanjin exercise could decrease depression, fatigue, anger and tension symptoms measured by SCL-90 scale substantially, and improve self-respect,

energy and emotional state after a 6-month intervention [53]. In the present trial, self-symptom intensity score in Baduanjin exercise group had a mean reduction of 7.3 points, and a mean reduction of 5.96 points in control group after 12 week exercise, but the difference was not significant between groups. Other psychological outcomes, such as stress, self-efficacy, self-esteem, mood and mindfulness, quality of life, and quality of sleep, were not different between groups either. We thought that the original activities habit practiced by the included participants, such as basketball, badminton, and ping-pong sport, might have a similar effect on psychological outcomes of college students as Baduanjin exercise. However, we did find the attention measured by the Schulte Grid test was improved with a mean shorten 7.5 second in participants over a 12 weeks of regular Baduanjin exercise comparing with controls ($P = 0.034$). This finding may be due to the fact that the exercisers were required to concentrate on adjusting breathing to make unity of mind and breathing when Baduanjin exercise was practicing.

Study strengths and limitations

This trial focused on a unique college student population and we used a rigorous randomized, parallel-controlled design with a relatively large sample to evaluate the effectiveness and safety of Baduanjin exercise on physical and mental health. We arranged two qualified physical teachers serving as the Baduanjin exercise instructors in order to ensure the standardization of exercise training for participants. Participants in Baduanjin exercise group were gathered to do the exercise at a fixed setting and time. To control confounding factors, all participants were required to record their daily physical activity. Furthermore, the result evaluators and statistical analysts were blinded to ensure the authenticity and objectivity of the trial results. We conducted this trial according to the protocol strictly to control the quality. Therefore the internal validity of the results in this trial was achieved.

Several limitations occurred in this trial. Ideally, everyone involved in an RCT should be blinded, but this is not always feasible in the non-pharmacological trials [54]. Although the participants and exercise instructors are not blinded and the psychological outcome measures are self-report, the assessment of the related physical fitness outcomes, lumbar muscle strength and lower limb proprioception function, and the

statistical analyses were performed by research staffs blinded to the treatment allocation. Second, all participants were recruited in one medical university with a greater proportion of female participants, and not controlled intensity, frequency and types of usual activities in two groups, and a relative inadequacy of adherence rate in Baduanjin exercise group. 18 participants had 75% attendance rate or less, and only 40 participants attained more than 85% attendance rate in Baduanjin exercise group. Third, the physical activity was self-report that has inherent limitations such as recall bias or social desirability. In addition, because the effect size between BDJ and Con groups was lower than anticipated, it was possible the study power was overestimated. All of which may affect the efficacy of Baduanjin intervention. Nonetheless, excellent protocol adherence, strictly performance supervision, no significant loss to follow-up and withdrawn, and the use of adjusted analyses strengthen the findings.

[Go to:](#)

Summary

In conclusion, this study demonstrates that regular Baduanjin exercise may improve lower limb proprioception function and cardiorespiratory endurance, enhance flexibility, explosive force of lower limb as well as attention in a college student population. Our finding indicates that regular Baduanjin exercise may be an effective, safe and useful form of exercise to promote the college students' physical health and attention.

[Go to:](#)

Differences Between Protocol and Trials

- 1 list-behavior=enumerated prefix-word= mark-type=decimal
- 2 Data of extension and lateroflexion lumbar muscle strength in all participants at baseline was lost due to accidental fault of the measurement machine (the *Tergumed Work Station*) at baseline assessment completed. They were therefore not involved in the final statistical analysis.
- 3 Through expert consultation, lower limb proprioception function was thought to be a better indicator to reflect the effect of Baduanjin exercise on proprioception function. As a result, lumbar proprioception function was replaced by lower limb proprioception function.

- 4 Standing long jump was discussed to be added in the secondary outcomes because it can excellently reflect explosive force of lower limb and also is an important component of physical fitness.

[Go to:](#)

Supporting Information

/article/body/sec/

S1 CONSORT Checklist

caption a9**CONSORT Checklist.**

(DOC)

[Click here for additional data file.](#)(226K, doc)

S1 Protocol

caption a9**Protocol of this trial (Chinese version).**

(DOC)

[Click here for additional data file.](#)(53K, doc)

S2 Protocol

caption a9**Protocol of this trial (English version).**

(DOC)

[Click here for additional data file.](#)(46K, doc)

[Go to:](#)

Acknowledgments

The authors thank Dr Xuemei Sui (University of South Carolina) for her helpful evaluation and revision on the final manuscript. This study is supported by Rehabilitation Technology of Collaborative Innovation Center, FJTCM (Grant No. X2012002-Collaboration), and the Special Scientific Research Fund of Public Welfare Profession of China (Grant No. 201307004), Ministry of Science and Technology and Ministry of Finance of the People's Republic of China.

[Go to:](#)

Funding Statement

This study is funded by the Rehabilitation Technology of Collaborative Innovation Center, FJTCM (Grant No. X2012002-Collaboration), the Special Scientific Research Fund of Public Welfare Profession of China (Grant No. 201307004), Ministry of Science and Technology (<http://www.most.gov.cn>) and Ministry of Finance of the People's Republic of China (<http://www.mof.gov.cn>). The funders had no role in study design, data collection and analysis, decision to publish, or preparation

of the manuscript.

Go to:

Data Availability

All relevant data are within the paper.

Go to:

References

1. Lepp A, Barkley JE, Sanders GJ, Rebold M, Gates P. The relationship between cell phone use, physical and sedentary activity, and cardiorespiratory fitness in a sample of U.S. college students. *Int J Behav Nutr Phys Act.* 2013; 10:79 doi: [10.1186/1479-5868-10-79](https://doi.org/10.1186/1479-5868-10-79) [[PMC free article](#)] [[PubMed](#)]
2. Owen N, Healy GN, Matthews CE, Dunstan DW. Too much sitting: the population health science of sedentary behavior. *Exerc Sport Sci Rev.* 2010;38:105–113. doi: [10.1097/JES.0b013e3181e373a2](https://doi.org/10.1097/JES.0b013e3181e373a2) [[PMC free article](#)] [[PubMed](#)]
3. Teiji N. The integration of school nutrition program into health promotion and prevention of lifestyle-related diseases in Japan. *Asia Pac J Clin Nutr.* 2008; 17: 349–351. [[PubMed](#)]
4. American College Health Association. American College Health Association National College Health Assessment (ACHA-NCHA) Spring 2005 Reference Group Data Report. *J Am Coll Health.* 2006;55:5–16. [[PubMed](#)]
5. Werch (Chad) CE, Bian H, Moore MJ, Ames S, DiClemente CC, Weiler RM. Brief multiple behavior interventions in a college student health care clinic. *J Adolesc Health.* 2007;41:577–585. [[PMC free article](#)] [[PubMed](#)]
6. National College Health Assessment Web Summary. American College Health Association. October 17, 2002. Available: http://www.acha.org/projects_programs/ncha_sampledata_public.cfm.
7. Nelson TF, Gortmaker SL, Subramanian SV, Wechsler H. Vigorous physical activity among college students in the United States. *J Phys Act Health.* 2007;4:495–508. [[PubMed](#)]
8. Hamilton MT, Hamilton DG, Zderic TW. Role of low energy expenditure and sitting in obesity, metabolic syndrome, type 2 diabetes, and cardiovascular disease. *Diabetes.* 2007; 56:2655–2667. [[PubMed](#)]
9. Matthews CE, Chen KY, Freedson PS, Buchowski MS, Beech BM,

- Pate RR, et al. Amount of time spent in sedentary behaviors in the United States, 2003–2004. *Am J Epidemiol*. 2008; 167:875–881. doi: [10.1093/aje/kwm390](https://doi.org/10.1093/aje/kwm390) [PMC free article] [PubMed]
10. Morrell JS, Cook SB, Carey GB. Cardiovascular fitness, activity, and metabolic syndrome among college men and women. *Metab Syndr Relat Disord*. 2013;11:370–376. doi: [10.1089/met.2013.0011](https://doi.org/10.1089/met.2013.0011) [PubMed]
11. Kang J, Ciecierski CC, Malin EL, Carroll AJ, Gidea M, Craft LL, et al. A latent class analysis of cancer risk behaviors among U.S. college students. *Prev Med*. 2014; 64:121–125. doi: [10.1016/j.ypmed.2014.03.023](https://doi.org/10.1016/j.ypmed.2014.03.023) [PMC free article] [PubMed]
12. St Kim, Kathy A, Anthony T. A systematic review: Students with mental health problems: A growing problem. *Int J of Nurs Pract*. 2010;16:1–6. [PubMed]
13. Stallman HM. Prevalence of psychological distress in university students—implications for service delivery. *Aust Fam Physician*. 2008;37:673–677. [PubMed]
14. Gore FM, Bloem PJ, Patton GC, Ferguson J, Joseph V, Coffey C, et al. Global burden of disease in young people aged 10–24 years: a systematic analysis. *Lancet*. 2011; 377:2093–2012. doi: [10.1016/S0140-6736\(11\)60512-6](https://doi.org/10.1016/S0140-6736(11)60512-6) [PubMed]
15. World Health Organization: *Global Burden of Disease: 2004 Update*. Geneva: WHO; 2008.
16. Dyrbye L, Thomas M, Shanafelt T. Systematic review of depression, anxiety, and other indicators of psychological distress among U.S. and Canadian medical students. *Acad Med*. 2006;81:354–373. [PubMed]
17. Hunt J, Eisenberg D. Mental health problems and help-seeking behavior among college students. *J Adolesc Health*. 2010, 46:3–10. doi: [10.1016/j.jadohealth.2009.08.008](https://doi.org/10.1016/j.jadohealth.2009.08.008) [PubMed]
18. Sagatun A, Sogaard AJ, Bjertness E, Selmer R, Heyerdahl S. The association between weekly hours of physical activity and mental health: A three-year follow-up study of 15-16-year-old students in the city of Oslo, Norway. *Bmc Public Health*. 2007;7 doi: [10.1186/1471-2458-7-155](https://doi.org/10.1186/1471-2458-7-155) [PMC free article] [PubMed]
19. Biddle SJ, Asare M. Physical activity and mental health in children and adolescents: a review of reviews. *Br J Sports Med*. 2011;45:886–895. doi: [10.1136/bjsports-2011-090185](https://doi.org/10.1136/bjsports-2011-090185) [PubMed]

20. Paluska SA, Schwenk TL. Physical activity and mental health: current concepts. *Sports Med.* 2000; 29:167–180. [[PubMed](#)]
21. Fox KR. The influence of physical activity on mental well-being. *Public Health Nutr.* 1999;2:411–418. [[PubMed](#)]
22. Miller KE, Hoffman JH. Mental Well-Being and Sport-Related Identities in College Students. *Sociol Sport J.* 2009;26:335–356. [[PMC free article](#)] [[PubMed](#)]
23. French SU, Alan LS, Cox AE. Attachment relationships and physical activity motivation of college students. *Psychology & Health.* 2011;26:1063–1081. [[PubMed](#)]
24. Van Dyck D, Teychenne M, McNaughton SA, De Bourdeaudhuij I, Salmon J. Relationship of the Perceived Social and Physical Environment with Mental Health-Related Quality of Life in Middle-Aged and Older Adults: Mediating Effects of Physical Activity. *PLoS ONE.* 2015;10: e0120475 doi: [10.1371/journal.pone.0120475](https://doi.org/10.1371/journal.pone.0120475) [[PMC free article](#)] [[PubMed](#)]
25. Yamazaki F, Yamada H, Moridawa S. Influence of an 8-week exercise intervention on body composition, physical fitness, and mental health in female nursing students. *J UOEH.* 2013; 35:51–58. [[PubMed](#)]
26. Ha CH, So WY. Effects of combined exercise training on body composition and metabolic syndrome factors. *Iran J Public Health.* 2012; 41:20–26. [[PMC free article](#)] [[PubMed](#)]
27. Chen HH, Yeh ML, Lee FY. The effects of Baduanjin qigong in the prevention of bone loss for middle-aged women. *Am J Chin Med.* 2006;34:741–747. [[PubMed](#)]
28. Koh TC. Baduanjin—an ancient Chinese exercise. *Am J Chin Med.* 1982, 10:14–21. [[PubMed](#)]
29. Mei L, Chen Q, Ge L, Zheng G, Chen J. Systematic review of Chinese traditional exercise baduanjin modulating the blood lipid metabolism. *Evid Based Complement Alternat Med.* 2012; 2012:282131 doi: [10.1155/2012/282131](https://doi.org/10.1155/2012/282131) [[PMC free article](#)] [[PubMed](#)]
30. An BC, Wang Y, Jiang X, Lu HS, Fang ZY, Wang Y, Dai KR. Effects of Baduanjin exercise on knee osteoarthritis: a one-year study. *Chin J Integr Med.* 2013;19:143–148. doi: [10.1007/s11655-012-1211-y](https://doi.org/10.1007/s11655-012-1211-y) [[PubMed](#)]
31. Hsu MC, Wang TS, Liu YP, Liu CF. Effects of Baduanjin exercise

on oxidative stress and antioxidant status and improving quality of life among middle-aged women. *Am J Chin Med.* 2008; 36:815–826.

[[PubMed](#)]

32. Liu HF, Zhao LM, An HY. A study of Baduanjin exercise for the mental health of college students. *J Beijing Sport Univ.* 2007; Suppl: 149–151.

33. Liu HF, An HY, Wang CH, Cao C, Zhang B. Psychological effects of health Qigong Baduanjin. *J Wuhan Inst of Phys Educ.* 2008; 42:54–58.

34. Xue ZH. A research on the effect of Baduanjin exercise on female university student's physique health. *J Anhui Sport Sci.* 2012; 33:46–49.

35. Zheng G, Li M, Lan X, Yan X, Lin Q, Chen L, et al. The effect of Baduanjin exercise for physical and psychological wellbeing of college students: study protocol for a randomized controlled trial. *Trials.* 2013;14:422 doi: [10.1186/1745-6215-14-422](https://doi.org/10.1186/1745-6215-14-422) [[PMC free article](#)]

[[PubMed](#)]

36. Health Qigong Management Center of General Administration of Sport of China: *Health qigong—Baduanjin*. Beijing: People's Sports Publishing House of China; 2003.

37. Wang ZY. SCL-90. *Shanghai Archives of Psychiatry.* 1984; 2:68–70.

38. Yang TZ, Huang HT. An epidemiological study of psychological pressure of urban residents in social transformation. *J Chin Epidemiol.* 2003; 24:760–764. [[PubMed](#)]

39. Zhang JX, Schwarzer R. Measuring optimistic self-beliefs: A Chinese adaptation of the General Self-efficacy Scale. *Psychologia.* 1995; 38:174–181.

40. Yang HG, Wei Z. Comment on evolution process of Lishi Baduanjin. *Journal of Xian Physical Education University.* 2013; 30:315–320.

41. Qu XF. An experimental research of how the core strength training influencing waist and abdominal muscle force and endurance of male undergraduates in common colleges and universities. *Liaoning Sport Science and Technology.* 2014;36:45–46.

42. Li S, Ala M, Wang L, Wu Y. The effects of progressive resistance exercise training on muscle strength of college students. *Journal of Jilin*

Institute of Physical Education. 2012;28:13–15.

43. Zhou XQ, Zeng YQ, Yang BL, Wang AL. Effects of health Qigong and Baduanjin on the middle and old aged people's blood lipid. *J Beijing Sport Univ.* 2007; 30:795–797.

44. Lin Q. Baduanjin and Chinese Medical Health Preservation. *J Fujian Univ Tradit Chin Med.* 2010; 20:55–56.

45. Jiang X. A shallow discussion on the dialectical relationship between Baduanjin and health keeping in Chinese medicine. *Chinese Wushu Research.* 2012;1:95–96.

46. Geng YQ, Wang XD. Effect of eight-length brocade exercise on mental sub-health regulation based on the theory of emotion controlled by five zang-organs. *China Journal of Tadtional Chinese Medicine and Pharmacy.* 2008;23:348–350.

47. Zhang X, Ren SJ. The effect of Baduanjin on lumbar muscle strain. *Chin Med Mod Distance Educ China.* 2011; 9:105.

48. Chen MC, Liu HE, Huang HY, Chiou AF. The effect of a simple traditional exercise programme (Baduanjin exercise) on sleep quality of older adults: a randomized controlled trial. *Int J Nurs Stud.* 2012; 49:265–273. doi: [10.1016/j.ijnurstu.2011.09.009](https://doi.org/10.1016/j.ijnurstu.2011.09.009) [[PubMed](#)]

49. Peng LY, Chen Y, Yao X. Effect of Baduanjin exercise training for physiological functions in college students. *Sport Research.* 2012; 2:115–116.

50. Yang HJ. Study of Baduanjin exercise for cardiopulmonary function in college students. *Sport Research and Education.* 2011;26:198–200.

51. Yu GB. *Chinese Qigong illustrated.* Beijing: New World Press, 1995.

52. Wang Y. The influence of health Qigong Baduanjin training on the psychological health of college students. *Journal of Beijing Sport University.* 2011; 34:102–104.

53. Mo GN. Experimental study of Baduanjin on the mental health of female university students. *Wushu Science.* 2013; 10:81–83.

54. Boutron I, Moher D, Altman DG, Schulz KF, Ravaud P. CONSORT Group: Extending the CONSORT statement to randomized trials of nonpharmacologic treatment: explanation and elaboration. *Ann Intern Med.* 2008; 148:295–309. [[PubMed](#)]